

## MENINGOCOCCUS

cannot afford to await culture results, and counter-current immunoelectrophoresis, although completed in a few minutes, is not widely available. A 15 to 30 percent mortality rate in the civilian population is too high and most likely results from the delay in therapy.

By keeping meningococcal disease in mind when confronted with a febrile patient in a toxic condition, particularly when a skin rash is noted, and instituting early therapy, you can increase the patient's chances for survival.

### REFERENCES

1. Gold R, Winklehake JL, Mars RS, et al: Identification of an epidemic strain of group C *Neisseria meningitidis* by bactericidal serotyping. *J Infect Dis* 124:593-597, Dec 1971
2. Counts GW, Steeley L, Beaty H: Identification of an epidemic strain of *Neisseria meningitidis* by bacteriocin typing. *J Infect Dis* 124:26-32, Jul 1971
3. Greenfield S, Shee PR, Feldman HA: Meningococcal carriage in a population of "normal" families. *J Infect Dis* 123:67-73, 1971
4. Goldschneider I, Gotschlich EC, Artenstein MS: Human immunity to the meningococcus—I. The role of humoral antibodies. *J Exp Med* 129:1307-1326, 1969
5. Frasch CE, Chapman SS: Classification of *Neisseria meningitidis* group B into distinct serotype—III. Application of a new bactericidal inhibition technique to distribution of serotypes among cases and carriers. *J Infect Dis* 127:149-154, 1973
6. Young LS, LaForce FM, Head JJ, et al: A simultaneous outbreak of meningococcal and influenza infections. *N Engl J Med* 287:5-8, Jul 6, 1972
7. Artenstein MS, Rust JH, Hunter DH, et al: Acute respiratory disease and meningococcal infection in army recruits. *JAMA* 201:1004-1008, Sep 25, 1967
8. Koppes GM, Ellenbogen C, Gebhart, RJ: Group Y meningococcal disease in United States Air Force recruits. *Am J Med* 62:661, 1977
9. Toews WH, Bass JW: Skin manifestations of meningococcal infection. *Am J Dis Child* 127:173-176, 1974
10. Davis CE, Arnold K: Role of meningococcal endotoxin in meningococcal purpura. *J Exp Med* 140:159-171, 1974
11. Fass RJ, Saslaw S: Chronic meningococcemia. *Arch Int Med* 130:943-946, 1972
12. McLean S, Caffey J: Endemic purpuric meningococcus bacteremia in early life. *Am J Dis Child* 42:1053-1074, 1931
13. Tompkins VN: The diagnostic value of smears taken from purpuric lesions of the skin in meningococcal disease. *JAMA* 123:31-32, 1943
14. Whittle HC, Greenwood BM, Davidson NM, et al: Meningococcal antigen in diagnosis and treatment of group A meningococcal infections. *Am J Med* 58:823-828, Jun 1975
15. Whittle HC, Abdullah MT, Fakunle FA, et al: Allergic complications of meningococcal disease. *Br Med J* 2:733-740, 1973
16. Blackfan KD: The treatment of meningococcal meningitis. *Medicine* 1:139-212, 1922
17. Souther PM, Sanford JP: Meningococcal meningitis—Suboptimal response to cephalothin therapy. *N Engl J Med* 280:1163-1164, 1969
18. Hardman JM, Earle KM: Myocarditis in 200 fatal meningococcal infections. *Arch Path* 87:318-325, 1969
19. Beal LR, Ustach TJ, Forker AD: Meningococcemia without meningitis presenting as cardiac tamponade. *Am J Med* 51:659-662, 1971
20. The Meningococcal Disease Surveillance Group: Meningococcal disease: Secondary attack rate and chemoprophylaxis in the United States, 1974. *JAMA* 235:261-265, 1976
21. Kuhns DM, Nelson CT, Feldman HA: The prophylactic value of sulfadiazine. *JAMA* 123:335-339, 1943
22. Guttler RB, Counts GW, Avent CK, et al: Effect of rifampin and minocycline on meningococcal carrier rates. *J Infect Dis* 124:199-205, 1971
23. Jacobson JA, Daniel B: Vestibular reactions associated with minocycline-antimicrobial agents and chemotherapy. 8:453-456, 1975
24. Gotschlich EC, Goldschneider I, Artenstein MS: Human immunity to the meningococcus—IV. Immunogenicity of group A and group C meningococcal polysaccharides in human volunteers. *J Exp Med* 129:1367-1384, 1969
25. Artenstein MS, Gold R, Zimmerly JG: Prevention of meningococcal disease by group C polysaccharide vaccine. *N Engl J Med* 282:417-420, 1970
26. Wyle FA, Artenstein MS, Brandt EC, et al: Immunologic response of man to group B meningococcal polysaccharide vaccines. *J Infect Dis* 126:514-522, 1972

## Early Diagnostic Esophagoscopy in the Management of Corrosive Burns of the Esophagus

AS EARLY AS WE CAN—it may be 10, 12, 14 or even 24 hours after injury—we do an esophagoscopy in all cases of possible corrosive burns of the esophagus. It's diagnostic esophagoscopy, because all we want to find out is if the esophagus is burned or if it's not burned. If the esophagus is not burned, a patient will be in hospital for a very short time for the treatment of the burns of the lips and buccal mucosa. If the esophagus is burned, we will continue therapy with antibiotics and steroids. Barium swallows do not help us because there could be severe damage, and the scarring and stricture formation takes usually weeks to occur. So, the mistake has been made to get a barium swallow at say two weeks after injury. The results may look pretty good, yet these patients have difficulty eating. In some of them there may even be aspiration of food and we wonder why they keep having these problems. So, I think diagnostic esophagoscopy is vital to see which patient you have to treat.

—GILBERT S. CAMPBELL, MD, Little Rock

Extracted from *Audio-Digest Surgery*, Vol. 24, No. 15, in the Audio-Digest Foundation's subscription series of tape-recorded programs. For subscription information: 1577 East Chevy Chase Drive, Glendale, CA 91206.